



# Number Theory

Book 2

"Trick" (or technique) for identifying which small integers have the form  $x^2 + y^2$  or  $x^2 + 3y^2$  or ...

Define  $\theta(t) = \sum_{n=-\infty}^{\infty} t^n = 1 + 2t + 2t^4 + 2t^9 + 2t^{16} + 2t^{25} + \dots = \sum_{n \in \mathbb{Z}} t^{n^2}$

$$\theta(t)^2 = (1 + 2t + 2t^4 + 2t^9 + 2t^{16} + 2t^{25} + \dots)(1 + 2t + 2t^4 + 2t^9 + 2t^{16} + 2t^{25} + \dots) = 1 + 4t + 4t^2 + 4t^4 + 8t^5 + 4t^8 + \dots + 12t^{25} + \dots$$

The powers of  $t$  appearing in this expansion are precisely the exponents expressible as a sum of two squares. The coefficient of  $t^n$  on the right hand side is the number of solutions of  $n = x^2 + y^2$  ( $x, y \in \mathbb{Z}$ )

eg.  $25 = x^2 + y^2$  has 12 solutions  $(\pm 5, 0), (0, \pm 5), (\pm 3, \pm 4), (\pm 4, \pm 3)$

$5 = x^2 + y^2$  has 8 solutions  $(\pm 2, \pm 1), (\pm 1, \pm 2)$

$6 = x^2 + y^2$  has 0 solutions.

$$\theta(t)^3 = 1 + 6t + 12t^2 + 8t^3 + 6t^4 + 24t^5 + 24t^6 + 12t^8 + \dots$$

$1 = x^2 + y^2 + z^2$  has six solutions  $(\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)$

$2 = \dots$  twelve  $(\pm 1, \pm 1, 0), (\pm 1, 0, \pm 1), (0, \pm 1, \pm 1)$

$7 = x^2 + y^2 + z^2$  has 0 solutions

$\theta(t)^4$  has positive coefficient of  $t^n$  for every positive integer  $n$

Lagrange's Theorem: every positive integer is a sum of four squares

Fundamental Theorem of Arithmetic: Every positive integer is uniquely expressible as a product of primes. We'll explain exactly what this says and we'll prove it.

Fundamental Theorem of Calculus

.. .. of Linear Algebra

.. .. of Algebra

FTA = Fundamental Theorem of Arithmetic:

Positive integers factor uniquely as products of primes.

eg.  $12 = 6 \times 2 = 2 \times 3 \times 2$

$12 = 3 \times 4 = 3 \times 2 \times 2$

$12 = (-2) \times (-6) = 2 \times 6$

$12 = \boxed{1} \times 12$

Ignore factors of  $\pm 1$  in factorization: these are units. (invertible elements). In  $\mathbb{Z}$ , the only units are  $\pm 1$ .

eg.  $\mathbb{Z}[i] = \{a+bi : a, b \in \mathbb{Z}\}$  (commutative ring with identity 1) we again have unique factorization. The units are  $\pm 1, \pm i$ .

$i \cdot (-i) = 1$

$1 \cdot 1 = 1$

$(-i) \cdot (-1) = 1$

$\mathbb{Z}[i]$  = "Gaussian integers"

$12 = 2 \times 2 \times 3 = \underbrace{(1+i)(1-i)(1+i)(1-i)}_3 = (i-1)(-i-1)(1+i)(1-i) 3$

Cannot be factored any further; they are irreducible.

$2+i = i(1-2i)$

- Elements of  $\mathbb{R}$ :
- zero
  - units
  - irreducible
  - reducible.

$10 = 2 \times 5 = (1+i)(1-i)(1+2i)(1-2i) = (1+i)(1-i)(2+i)(2-i)$

$\times i$

$\times (-i)$

"migration of units"

A (commutative ring with identity) has unique factorization if every nonzero element can be factored as a product of irreducible elements and this factorization is essentially unique (ie. unique up to permutation of factors and migration of units).

Why don't we just say primes and composites instead of irreducible and reducible elements?

In  $\mathbb{Z}$ :  
zero: 0  
units:  $\pm 1$   
irreducible:  $\pm 2, \pm 3, \pm 5, \pm 7, \pm 11, \dots$   
reducible:  $\pm 4, \pm 6, \pm 8, \pm 9, \pm 10, \pm 12, \dots$

In  $\mathbb{Z}[i]$ :  
zero: 0  
units:  $\pm 1, \pm i$   
irreducibles:  $\pm 3 = \{1+i, 1-i, -1+i, -1-i\}$   
                  3           actually 3, -3, 3i, -3i  
 $\pm 1 \pm 2i, \pm 2i$   
etc.  
reducibles: 2 (actually 2, -2, 2i, -2i)

A nonzero element  $\alpha \in R$  is reducible if  $\alpha \neq 0$ ,  $\alpha \neq \text{unit}$  and  $\alpha = \alpha_1 \alpha_2$  with  $\alpha_1, \alpha_2$  not units.

$\alpha$  is irreducible if the only factorizations  $\alpha = \alpha_1 \alpha_2$  have either  $\alpha_1$  or  $\alpha_2$  is a unit.

eg. in  $\mathbb{Z}[i]$ , 3 is irreducible. If  $3 = \beta \gamma$ ,  $\beta, \gamma \in \mathbb{Z}[i]$  then either  $\beta \in \{\pm 1, \pm i\}$  or  $\gamma \in \{\pm 1, \pm i\}$ .  
2 is reducible since  $2 = (1+i)(1-i)$ , neither  $1+i$  nor  $1-i$  is a unit.

Eg. the ring  $\mathbb{Z}[\sqrt{5}] = \{a + b\sqrt{5} : a, b \in \mathbb{Z}\}$  does not have unique factorization.

$$\mathbb{Z}[\sqrt{6}] = \{a + b\sqrt{6} : a, b \in \mathbb{Z}\}$$

An example of non-unique factorization in  $\mathbb{Z}[\sqrt{6}]$ :  $6 = 2 \times 3 = \sqrt{6} \times \sqrt{6}$   
 $2, 3, \sqrt{6}$  are irreducible in  $\mathbb{Z}[\sqrt{6}]$ .

But be careful:  $\mathbb{Z}[\sqrt{6}]$  has infinitely many units.

Pell's equation  $x^2 - 6y^2 = 1$  has solutions  $(\pm 1, 0), (\pm 5, \pm 2)$

$x^2 - 6y^2 = -1$  has no integer solutions.

(Look at the equation mod 3)

$$(5 + 2\sqrt{6})(5 - 2\sqrt{6}) = 1 \quad \text{in } \mathbb{Z}[\sqrt{6}]$$

In  $\mathbb{Z}[\sqrt{5}]$ , the only units are  $\pm 1$ .

Unique factorization is not universal.

$6 = 2 \times 3 = (1 + \sqrt{5})(1 - \sqrt{5})$   
where all factors  $2, 3, 1 + \sqrt{5}, 1 - \sqrt{5}$  are irreducible in  $\mathbb{Z}[\sqrt{5}]$   
It holds in  $\mathbb{Z}, \mathbb{Z}[i], \mathbb{Z}[\sqrt{2}], \mathbb{Z}[\sqrt{3}], \mathbb{Z}[\sqrt{5}], \mathbb{Z}[\sqrt{2}], \mathbb{Z}[\sqrt{3}]$   
Not in  $\mathbb{Z}[\sqrt{5}]$  or  $\mathbb{Z}[i]$ .